

Alistair Savage


Fields Institute and University of Toronto

**Quiver Varieties  
and  
Geometric Representation Theory**

---

## Geometric Representation Theory

---



---

## A Sample Lie Algebra: $\mathfrak{sl}_n$

---

$\mathfrak{sl}_n = n \times n$  traceless matrices

$$[A, B] = AB - BA$$

Presentation in terms of *Chevalley generators*

$$e_i = E_{i,i+1}, \quad f_i = E_{i+1,i}, \quad i = 1, \dots, n-1$$

and certain relations.

A *representation* of  $\mathfrak{sl}_n$  is

- A complex vector space  $V$
- A map  $\mathfrak{sl}_n \rightarrow \text{End}(V)$  such that

$$[A, B](v) = A(B(v)) - B(A(v))$$

---

**Dictionary**

---

simply-laced Kac-Moody algebra  $\leftrightarrow \mathfrak{sl}_n$

irreducible integrable highest weight rep  $\leftrightarrow$  f.d. rep

weight space of a rep  $\leftrightarrow$  eigenspace of diagonal matrices

---

## Quiver Variety Approach

---

$\mathfrak{g}$  = simply-laced Kac-Moody Lie algebra (e.g.  $\mathfrak{g} = \mathfrak{sl}_n$ )

$L$  = irreducible integrable highest weight rep of  $\mathfrak{g}$

$L$  decomposes into weight spaces:

$$L = \bigoplus_{\lambda} L_{\lambda}$$

$\lambda \longleftrightarrow$  quiver variety  $QV(\lambda)$ .

---

## Quiver Variety Approach

---

Weight space  $L_\lambda \longleftrightarrow$  “Homology” of  $QV(\lambda)$

$\dim L_\lambda \longleftrightarrow$  # irr. comps. of  $QV(\lambda)$

Action of  $\mathfrak{g} \longleftrightarrow$  Correspondences

### Correspondences:

$QV(\lambda_1) \leftarrow$  “Intermediate Variety”  $\rightarrow QV(\lambda_2)$

---

## Quivers

---


quiver = oriented graph

### Examples:

1. Quiver of type  $A_n$



2. Quiver of type  $A_n^{(1)}$



---

## Representations of Quivers


---

### Representation of a quiver:

vertex  $\longrightarrow$  f.d. vector space

arrow  $\longrightarrow$  linear map

**Example:** A representation of the quiver



consists of

$V_1, V_2$  - f.d.  $\mathbb{C}$ -vector spaces

$x \in \text{Hom}(V_2, V_1)$



---

## Quiver Varieties


---

$\mathfrak{g}$  = simply-laced Kac-Moody Lie algebra (e.g.  $\mathfrak{g} = \mathfrak{sl}_{n+1}$ )


Consider the Dynkin graph of  $\mathfrak{g}$



Take *both* orientations of each edge



Add in *shadow vertices*



Call this quiver  $Q(\mathfrak{g})$

---

## Quiver Varieties

---


$I =$  set of vertices of Dynkin graph of  $\mathfrak{g}$

Fix collections of v.s.  $\mathbf{V} = (V_k)_{k \in I}$ ,  $\mathbf{W} = (W_k)_{k \in I}$

Let  $\mathbf{v} = (\dim V_k)_{k \in I}$ ,  $\mathbf{w} = (\dim W_k)_{k \in I}$

Define (Nakajima)

$$\mathcal{L}(\mathbf{v}, \mathbf{w}) = \{ \text{reps of } Q(\mathfrak{g}) \text{ with v.s. } \mathbf{V} \text{ and } \mathbf{W} \\ + \text{ conditions} \} / \prod_{k \in I} GL(V_k)$$



---

## Alternative Description

---

$\mathbf{M}(\mathbf{v}, \mathbf{w}) =$  space of reps of quiver with fixed vector spaces

$$\begin{array}{c} \mathbf{M}(\mathbf{v}, \mathbf{w}) //_{\xi_1} \prod_k GL(V_k) \longleftarrow \text{hyper-Kähler quotient (smooth)} \\ \pi \downarrow \\ \mathbf{M}(\mathbf{v}, \mathbf{w}) //_{\xi_2} \prod_k GL(V_k) \longleftarrow \text{hyper-Kähler quotient (singular)} \end{array}$$

$$\mathcal{L}(\mathbf{v}, \mathbf{w}) = \pi^{-1}(0)$$

$\mathcal{L}(\mathbf{v}, \mathbf{w})$  is a Lagrangian subvariety and a deformation retract of

$$\mathcal{M}(\mathbf{v}, \mathbf{w}) \stackrel{\text{def}}{=} \mathbf{M}(\mathbf{v}, \mathbf{w}) //_{\xi_1} \prod_k GL(V_k)$$

---

## Weights

---

Want to construct irred. integ. highest weight rep of  $\mathfrak{g}$ .

In  $\mathcal{L}(\mathbf{v}, \mathbf{w})$ ,

$\mathbf{w} \longleftrightarrow$  highest weight of rep

$$\mathbf{w} \longleftrightarrow \sum_{k \in I} w_k \Lambda_k$$

$\Lambda_k$  – fundamental weights

$\mathbf{v} \longleftrightarrow$  weight space

$$\mathbf{v} \longleftrightarrow \sum_{k \in I} w_k \Lambda_k - \sum_{k \in I} v_k \alpha_k$$

$\alpha_k$  – simple roots

---

**Example:  $\mathfrak{sl}_2$** 

---

Let  $\mathfrak{g} = \mathfrak{sl}_2$ .




$$\begin{aligned}\mathcal{L}(k, n) &= \{B \in \text{Hom}(\mathbb{C}^n, \mathbb{C}^k) \mid B \text{ surjective}\} / GL(\mathbb{C}^k) \\ &\cong Gr(n - k, n)\end{aligned}$$

---

**Example: Adjoint Representation of  $\mathfrak{sl}_3$** 


---

Adjoint rep of  $\mathfrak{sl}_3$  has h.w.  $\Lambda_1 + \Lambda_2 \Rightarrow w = (1, 1)$ .




---

## Other Examples of Quiver Varieties

---

- $\mathfrak{g} = \mathfrak{sl}_n$ , highest weight  $a\Lambda_1$   
→ partial flag varieties
- $\mathfrak{g}$  affine,  $w = 0$ ,  $v \leftrightarrow$  imaginary root  
→ ALE spaces (resolutions of simple singularities)
- $\mathfrak{g}$  affine  
→ moduli spaces of instantons on ALE spaces
- Jordan quiver



→ Hilbert schemes of points in  $\mathbb{C}^2$

---

## Benefits of Geometric Approach

---

- Alternative (often simpler) geometric proofs of algebraic facts
- Rep theory organizes homological information
- Connection to crystal graphs ( $q \rightarrow 0$  limit of quantum groups)
- Geometrically defined bases with remarkable properties



---

## Geometrically Defined Bases

---

Recall,

weight space  $\longleftrightarrow$  “homology” of  $QV$   
 dimension of weight space  $\longleftrightarrow$   $\#$  irr. comps. of  $QV$

Classes of irr. comps. of  $QV$  yield basis of representation

<b>Homology Theory</b>	<b>Basis</b>
Constructible functions	Semicanonical basis
Top dim Borel-Moore homology	Semicanonical basis?
Perverse sheaves	Canonical basis

Nice positivity, integrality, and compatibility properties

---

## Nice Properties of Geometric Bases

---

### Positivity & Integrality:

$$f_k \cdot b = \sum c_j b_j, \quad c_j \in \mathbb{Z}_{\geq 0}$$

### Compatibility:

Have geometric basis  $B$  of  $U^-(\mathfrak{g})$

For *any* irred hw rep  $L$  of  $\mathfrak{g}$  with hw vec  $v$ ,


$$\{b \cdot v \mid b \in B, b \cdot v \neq 0\}$$

is a basis of  $L$ .

---

## Geometric Boson-Fermion Correspondence

---



**Reference:** Savage, arXiv:math.RT/0508438

---

## Bosonic Fock Space

---

Infinite-dim Heisenberg algebra:

$$\mathfrak{s} = \bigoplus_{m \in \mathbb{Z} \setminus \{0\}} \mathbb{C}s_m \oplus \mathbb{C}K$$

$$[\mathfrak{s}, K] = 0, \quad [s_m, s_n] = m\delta_{m, -n}K$$

Bosonic Fock Space:

$$B = \mathbb{C}[p_1, p_2, \dots]$$

Action of  $\mathfrak{s}$  on  $B$  by

$$s_m \mapsto m \frac{\partial}{\partial p_m}, \quad s_{-m} \mapsto p_m, \quad m > 0$$

$$K \mapsto \text{Id}$$

---

## Fermionic Fock Space

---

$$F = \text{Span}_{\mathbb{C}}\{\underline{i}_0 \wedge \underline{i}_1 \wedge \underline{i}_2 \wedge \dots \mid i_k \in \mathbb{Z}, i_0 > i_1 > \dots, i_k = -k \text{ for } k \gg 0\}$$

= “infinite wedge space”

Part of a larger Fock space with action of an infinite Clifford algebra

$\mathfrak{gl}_{\infty}$  (and  $\mathfrak{sl}_{\infty}$ ) act on  $F$  by derivations:


For  $a \in \mathfrak{gl}_{\infty}$ ,

$$a(\underline{i}_0 \wedge \underline{i}_1 \wedge \dots) = (a \cdot \underline{i}_0) \wedge \underline{i}_1 \wedge \dots + \underline{i}_0 \wedge (a \cdot \underline{i}_1) \wedge \dots + \dots$$

---

## Boson-Fermion Correspondence

---



---

## Boson-Fermion Correspondence

---

Action of  $\mathfrak{s}$  on  $F$ : “Bosonization”

$$s_m \mapsto \sum_{j \in \mathbb{Z}} E_{j, j+m}, \quad m \in \mathbb{Z} \setminus \{0\},$$

$$K \mapsto \text{Id}$$

$$F \cong B \text{ as } \mathfrak{s}\text{-modules}$$


Action of Clifford algebra on  $B$  (generating functions, vertex operator algebras): “Fermionization”

$$F \cong B \text{ as Clifford algebra modules}$$

---

## Hilbert Schemes

---



$$X_n = \{(B_1, B_2, i) \mid B_j \in \text{End } V, i \in \text{Hom}(\mathbb{C}, V), + \text{ conditions}\} / GL(V)$$

Here,  $V \cong \mathbb{C}^n$ .

### Conditions:

- $B_1 B_2 = B_2 B_1$
- $\text{im } i$  generates  $V$  under  $B_1, B_2$



---

## Hilbert Schemes

---

$X_n \cong$  Hilbert scheme of  $n$  points in  $\mathbb{C}^2$

Hilbert scheme is a resolution of singularities:

$$X_n \rightarrow (\mathbb{C}^2)^n / S_n$$

Have  $T = \mathbb{C}^*$  action on  $X_n$ :

$$z \cdot (GL(V) \cdot (B_1, B_2, i)) = GL(V) \cdot (zB_1, z^{-1}B_2, i)$$

---

## Geometric Bosonic Fock Space

---

$$\mathbb{H}_n^B = H_T^{2n}(X_n), \quad \mathbb{H}^B = \bigoplus_{n=0}^{\infty} \mathbb{H}_n^B$$

**Correspondences:** Natural projections

$$X_{n+k} \xleftarrow{\pi_1} X_{n+k} \times X_n \xrightarrow{\pi_2} X_n$$

Define  $\mathfrak{p}_{-k} \in \text{End } \mathbb{H}^B$ ,  $k \geq 0$ , by

$$\mathfrak{p}_{-k}(\alpha) = (\pi_1)_!(\pi_2^* \alpha \cup [\Sigma_{n,k}]), \quad \alpha \in \mathbb{H}_n^B$$

Here

$$\begin{aligned} \Sigma_{n,k} &\subset X_{n+k} \times X_n \\ &\leftrightarrow \text{“adding } k \text{ points” at } z \in \mathbb{C} \times \{0\} \subset \mathbb{C}^2 \end{aligned}$$

---

## Geometric Bosonic Fock Space

---

Define adjoint operator  $\mathfrak{p}_k$ ,  $k > 0$

**Prop** (Nakajima, Vasserot):

$$[\mathfrak{p}_k, \mathfrak{p}_l] = k\delta_{k,-l}\text{Id}$$

So

$$s_k \mapsto \mathfrak{p}_k, \quad K \mapsto \text{Id}$$


defines action of  $\mathfrak{s}$  on  $\mathbb{H}^B$  and

$$\mathbb{H}^B \cong B \text{ as } \mathfrak{s}\text{-modules}$$

---

## Quiver Varieties for $\mathfrak{sl}_\infty$

---



$V =$  f.d.  $\mathbb{Z}$ -graded complex vector space

$$\mathbf{v} = \dim V = (\dim V_k)_{k \in I}$$

$$\mathbf{w} = \mathbf{e}_0$$

$$\mathcal{M}(\mathbf{v}, \mathbf{e}_0) = \{(B_1, B_2, i) \mid B_j \in \text{End } V, i \in \text{Hom}(\mathbb{C}, V_0) \\ + \text{conditions}\} / \prod_k GL(V_k)$$

### Conditions:

- $B_1 \in \text{End } V$ ,  $\deg B_1 = 1$  (i.e.  $B(V_k) \subseteq V_{k+1}$ )
- $B_2 \in \text{End } V$ ,  $\deg B_2 = -1$  (i.e.  $B(V_k) \subseteq V_{k-1}$ )
- $B_1 B_2 = B_2 B_1$
- $\text{im } i$  generates  $V$  under  $B_1, B_2$

---

## Geometric Fermionic Fock Space

---

$$\mathbb{H}^F = \bigoplus_{\mathbf{v}} H_T^{2|\mathbf{v}|}(\mathcal{M}(\mathbf{v}, \mathbf{e}_0)) \quad (\text{trivial } T\text{-action})$$

**Correspondences:** Have natural projections

$$\mathcal{M}(\mathbf{v} + \mathbf{e}_k, \mathbf{e}_0) \longleftarrow \mathcal{M}(\mathbf{v} + \mathbf{e}_k, \mathbf{e}_0) \times \mathcal{M}(\mathbf{v}, \mathbf{e}_0) \longrightarrow \mathcal{M}(\mathbf{v}, \mathbf{e}_0)$$

Define geometric action of  $\mathfrak{sl}_\infty$  (similar to Hilbert scheme picture) and

$$\mathbb{H}^F \cong F \text{ as } \mathfrak{sl}_\infty\text{-modules}$$

---

## Torus Fixed Points of Hilbert Schemes

---

At a  $T$ -fixed point of  $X_n$ , weight decomposition of  $V$  gives grading

$$V = \bigoplus_{m \in \mathbb{Z}} V_m$$

Can show that

- $B_1, B_2$  have degrees 1 and -1 resp.
- $\text{im } i \subseteq V_0$

Thus,

$T$ -fixed points of H.S. = quiver varieties for  $\mathfrak{sl}_\infty$

---

## Localization and the Boson-Fermion Correspondence

---

$X$  smooth with a  $T$ -action

$X^T = T$ -fixed points of  $X$

Localization theorem states

$$H_T^*(X) \otimes \mathbb{C}(t) \cong H_T^*(X^T) \otimes \mathbb{C}(t)$$

Thus (after a few technical steps),


$$\mathbb{H}^B = \bigoplus_n H_T^{2n}(X_n) \cong \bigoplus_n H_T^{2n}(X_n^T) = \mathbb{H}^F$$

A geometric boson-fermion correspondence!

---

## Further Directions

---



Fixed points of  $\mathbb{Z}/n\mathbb{Z} \subset T$  yield quiver varieties for  $\widehat{\mathfrak{sl}}_n$

Vertex operator construction of basic rep of  $\widehat{\mathfrak{sl}}_n$  should fit into this geometric picture

Should help give algebraic description of nice geometric bases



---

## Crystal Graphs

---

Quantum group  $U_q(\mathfrak{g})$  is a  $q$ -deformation of  $U(\mathfrak{g})$

Representations of  $\mathfrak{g}$  (or  $U(\mathfrak{g})$ ) have a  $q$ -deformation

usual rep theory  $\xleftarrow{q \rightarrow 1}$  quantum groups  $\xrightarrow{q \rightarrow 0}$  crystals

In  $q \rightarrow 0$  limit, rep theory becomes combinatorics


---

## Crystal graphs

---

$L = \text{rep of } U_q(\mathfrak{g})$

Fix a basis of  $L$  and depict elements of the basis by vertices:




---

## Crystal graphs

---

Consider the action of a Chevalley generator, say  $f_2$ :



“Nice” basis:

- At most one coeff of 1 in each expression
- All other coeffs have positive power of  $q$


---

## Crystal graphs

---

Take  $q \rightarrow 0$  limit

Label remaining edges by Chevalley generator index:




---

## Crystal graphs

---

Repeat for remaining Chevalley generators:



Obtain the *crystal graph* of rep  $L$

---

## Crystal Graphs

---

- Connected graph  $\Rightarrow$  irreducible rep
- Can compute characters by counting vertices of fixed weight
- Tensor product rule

“Nice” basis = canonical basis !!

**Problem:** How do we “thaw” a crystal?

---


## Tensor Product Rule

---

Let  $\mathfrak{g} = \mathfrak{sl}_2$ .

Consider  $V_3 \otimes V_4$ .

Crystal graphs are




---

 Tensor Product Rule
 

---

Tensor product rule yields



Thus

$$V_3 \otimes V_4 = V_2 \oplus V_4 \oplus V_6$$



---

## Realizations of Crystal Graphs

---

### Combinatorial Realizations:

- Young tableaux (classical Lie algebras)
- Young walls (affine Lie algebras)
- Kyoto path model (affine Lie algebras)
- Littelmann path model

### Geometric Realization:

Vertices of crystal graph = irred. comps. of QVs

Crystal operators (edges) defined geometrically

---

## Connections Between Realizations

---

*Finite and affine types A and D:* Explicit isomorphism between Young tableaux/wall realizations and geometric realization (S)


### Advantages:

- Explicit description of irr comps of QVs
- Gives geometric interpretation and suggests extensions of combinatorial constructions
- General QV theory gives universal method to “thaw” crystals

---

## Connections Between Realizations

---



---

## Other Constructions

---

### Extension to non-simply laced case:

- Crystal structure – done (S)
- “Full” structure – open

### Other constructions:

- Tensor products (Nakajima, Malkin)
- Fusion products (Schiffmann-S)
- Demazure modules (S)
- Spin representations, Clifford algebras (S)
- Virasoro algebra (Lehn)
- Others? Lie superalgebras? Jordan (super)algebras?

---

## Connections to Affine Grassmannian Approach

---

