

# Advances in Heisenberg categorification

$$\begin{array}{c} \nearrow \circ \searrow \\ \swarrow \nearrow \end{array} - \begin{array}{c} \nearrow \searrow \\ \swarrow \nearrow \circ \end{array} = \sum_{b \in B} \begin{array}{c} \uparrow \circ \\ \uparrow \circ \end{array} b^{\vee}$$

Alistair Savage  
University of Ottawa

Slides available online: [alistairsavage.ca/talks](http://alistairsavage.ca/talks)

Preprint: [arXiv:1802.01626](https://arxiv.org/abs/1802.01626)

# Outline

## Goals:

- 1 Describe a family of categories that categorify the Heisenberg algebra
- 2 Explain the relation to previous Heisenberg categories

## Overview:

- 1 Strict monoidal categories and string diagrams
- 2 The Frobenius Heisenberg category
- 3 Actions on categories of modules
- 4 Work in progress:  $q$ -deformations

# Strict monoidal categories

A **strict monoidal category** is a category  $\mathcal{C}$  equipped with

- a bifunctor (the **tensor product**)  $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ , and
- a **unit object**  $\mathbf{1}$ ,

such that

- $(A \otimes B) \otimes C = A \otimes (B \otimes C)$  for all objects  $A, B, C$ ,
- $\mathbf{1} \otimes A = A = A \otimes \mathbf{1}$  for all objects  $A$ .

## Remark: Non-strict monoidal categories

In a (not necessarily strict) **monoidal category**, the equalities above are replaced by isomorphism, and we impose some **coherence conditions**.

Every monoidal category is monoidally equivalent to a strict one.

## $\mathbb{k}$ -linear monoidal categories

Fix a commutative ground ring  $\mathbb{k}$ .

A **strict  $\mathbb{k}$ -linear monoidal category** is a strict monoidal category such that

- each morphism space is a  $\mathbb{k}$ -module,
- composition of morphisms is  $\mathbb{k}$ -bilinear,
- tensor product of morphisms is  $\mathbb{k}$ -bilinear.

### The interchange law

The axioms of a strict monoidal category imply the **interchange law**: For  $A_1 \xrightarrow{f} A_2$  and  $B_1 \xrightarrow{g} B_2$ , the following diagram commutes:

$$\begin{array}{ccc} A_1 \otimes B_1 & \xrightarrow{1 \otimes g} & A_1 \otimes B_2 \\ f \otimes 1 \downarrow & \searrow f \otimes g & \downarrow f \otimes 1 \\ A_2 \otimes B_1 & \xrightarrow{1 \otimes g} & A_2 \otimes B_2 \end{array}$$

## Categorification via split Grothendieck group

Suppose  $\mathcal{C}$  is an additive category (i.e. have  $\oplus$ ).

$\text{Iso}_{\mathbb{Z}}(\mathcal{C}) =$  free abelian group generated by isom. classes of objects in  $\mathcal{C}$ .

The **split Grothendieck group** of  $\mathcal{C}$  is

$$K_0(\mathcal{C}) = \text{Iso}_{\mathbb{Z}}(\mathcal{C}) / \langle [X \oplus Y] = [X] + [Y] \mid X, Y \in \mathcal{C} \rangle.$$

If  $\mathcal{C}$  is **monoidal**, then  $K_0(\mathcal{C})$  is a ring:

$$[X] \cdot [Y] = [X \otimes Y].$$

### Categorification

For our purposes, to **categorify** a ring  $R$  is to find an additive monoidal category  $\mathcal{C}$  such that

$$K_0(\mathcal{C}) \cong R \quad \text{as rings.}$$

# The Heisenberg algebra

Let  $\mathfrak{h}$  be the infinite-dimensional Heisenberg Lie algebra.

Thus,  $\mathfrak{h}$  is the complex Lie algebra with basis

$$\{c, q_n^\pm : n \geq 1\}$$

and product

$$[q_m^+, q_n^+] = [q_m^-, q_n^-] = [c, q_n^\pm] = 0, \quad [q_m^+, q_n^-] = \delta_{m,n} n c.$$

The associative Heisenberg algebra at **central charge**  $\xi \in \mathbb{Z}$  is

$$U(\mathfrak{h}) / \langle c - \xi \rangle.$$

We will describe categories that categorify these algebras.

# String diagrams

Fix a strict monoidal category  $\mathcal{C}$ .

We will denote a morphism  $f: A \rightarrow B$  by:




The **identity map**  $\text{id}_A: A \rightarrow A$  is a string with no label:




We sometimes omit the object labels when they are clear or unimportant.

# String diagrams


Composition is **vertical stacking** and tensor product is **horizontal juxtaposition**:



The **interchange law** then becomes:



A morphism  $f: A_1 \otimes A_2 \rightarrow B_1 \otimes B_2$  can be depicted:





# Presentations of strict monoidal categories

One can give **presentations** of some strict  $\mathbb{k}$ -linear monoidal categories, just as for monoids, groups, algebras, etc.

**Objects:** If the objects are generated by some collection  $A_i, i \in I$ , then we have all possible tensor products of these objects:

$$\mathbf{1}, \quad A_i, \quad A_i \otimes A_j \otimes A_k \otimes A_\ell, \quad \text{etc.}$$

**Morphisms:** If the morphisms are generated by some collection  $f_j, j \in J$ , then we have all possible compositions and tensor products of these morphisms (whenever these make sense):

$$\text{id}_{A_i}, \quad f_j \otimes (f_i f_k) \otimes (f_\ell), \quad \text{etc.}$$

We then often impose some **relations** on these morphism spaces.

**String diagrams:** We can build complex diagrams out of our simple generating diagrams.

# Monoidally generated symmetric groups

Define a strict  $\mathbb{k}$ -linear monoidal category  $\mathcal{S}$  with one generating object  $\uparrow$  and denote

$$\text{id}_{\uparrow} = \uparrow$$

We have one generating morphism

$$\begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} : \uparrow \otimes \uparrow \rightarrow \uparrow \otimes \uparrow.$$

We impose the relations:

$$\begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} = \begin{array}{c} \uparrow \\ \uparrow \end{array}, \quad \begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} = \begin{array}{c} \nearrow \\ \searrow \\ \searrow \\ \nearrow \end{array}.$$

Then

$$\text{End}_{\mathcal{S}}(\uparrow^{\otimes n}) = \mathbb{k}S_n$$

is the group algebra of the **symmetric group** on  $n$  letters.

# The degenerate affine Hecke category

Start again with the strict  $\mathbb{k}$ -linear monoidal category  $\mathcal{S}$ , but add a morphism:

$$\begin{array}{c} \uparrow \\ \circ \\ \uparrow \end{array} : \uparrow \rightarrow \uparrow$$

We impose the additional relation:

$$\begin{array}{c} \swarrow \circ \searrow \\ \nearrow \nwarrow \end{array} - \begin{array}{c} \swarrow \nwarrow \\ \nearrow \circ \searrow \end{array} = \begin{array}{c} \uparrow \\ \uparrow \end{array} .$$

Now

$$\text{End}(\uparrow^{\otimes n})$$

is the **degenerate affine Hecke algebra** (of type  $A$ ).

# The wreath product category

Fix an associative  $\mathbb{k}$ -algebra  $F$ . We add an endomorphism of  $\uparrow$  for each element of  $F$ .

More precisely, let  $\mathcal{W}(F)$  be the strict  $\mathbb{k}$ -linear monoidal category obtained from  $\mathcal{S}$  by adding morphisms such that we have an algebra homomorphism:

$$F \rightarrow \text{End } \uparrow, \quad f \mapsto \begin{array}{c} \uparrow \\ \bullet \\ f \end{array}$$

We impose the additional relations:

$$\begin{array}{c} \nearrow \\ \bullet \\ f \\ \searrow \end{array} = \begin{array}{c} \nearrow \\ f \\ \bullet \\ \searrow \end{array}, \quad f \in F$$

# The wreath product category

$$\text{End}_{\mathcal{W}(F)}(\uparrow^{\otimes n}) = F^{\otimes n} \rtimes S_n$$

is a wreath product algebra.

As a vector space,

$$F^{\otimes n} \rtimes S_n = F^{\otimes n} \otimes_{\mathbb{k}} \mathbb{k}S_n.$$

Multiplication is determined by

$$(f_1 \otimes \pi_1)(f_2 \otimes \pi_2) = f_1(\pi_1 \cdot f_2) \otimes \pi_1\pi_2, \quad f_1, f_2 \in F^{\otimes n}, \pi_1, \pi_2 \in S_n,$$

where  $\pi_1 \cdot f_2$  denotes the natural action of  $S_n$  on  $F^{\otimes n}$  by permutation of the factors.

**Note:**  $\mathcal{W}(\mathbb{k}) = \mathcal{S}$ , the symmetric group category.

**Want:** An affine version of the wreath product category.  $F = \mathbb{k}$  should recover the degenerate affine Hecke category.

# Frobenius algebras: Definition

## Frobenius algebra

A **Frobenius algebra** is a f.d. associative algebra  $F$  together with a linear trace map

$$\mathrm{tr}: F \rightarrow \mathbb{k}$$

such that the induced map

$$F \rightarrow \mathrm{Hom}_{\mathbb{k}}(F, \mathbb{k}), \quad f \mapsto (g \mapsto \mathrm{tr}(gf)),$$

is an isomorphism.

For simplicity, we assume that the trace is symmetric:

$$\mathrm{tr}(fg) = \mathrm{tr}(gf), \quad \text{for all } f, g \in F.$$

# Frobenius algebras: Examples

## Example ( $\mathbb{k}$ )

$\mathbb{k}$  is a Frobenius algebra with  $\text{tr} = \text{id}_{\mathbb{k}}$ .

## Example (Matrix algebra)

Any matrix algebra over a field is a Frobenius algebra with the usual trace.

## Example ( $\mathbb{k}[x]/(x^k)$ )

$\mathbb{k}[x]/(x^k)$  is a Frobenius algebra with

$$\text{tr}(x^\ell) = \delta_{\ell, k-1}.$$

# Frobenius algebras: Examples

## Example (Group algebra)

Suppose  $G$  is a finite group.

The **group algebra**  $\mathbb{k}G$  is a Frobenius algebra with

$$\mathrm{tr}(g) = \delta_{g,1_G}, \quad g \in G.$$

## Example (Zigzag algebra)

Associated to every quiver is a **zigzag algebra**. These are Frobenius algebras.

## Example (Hopf algebras)

Every f.d. Hopf algebra is a Frobenius algebra.

**From now on:**  $F$  is a Frobenius algebra with trace  $\mathrm{tr}$ .



# Frobenius algebras: dual bases

Fix a basis  $B$  of  $F$ . The dual basis is

$$B^\vee = \{b^\vee \mid b \in B\}$$

defined by

$$\mathrm{tr}(b^\vee c) = \delta_{b,c}, \quad b, c \in B.$$

It is easy to check that

$$\sum_{b \in B} b \otimes b^\vee \in F \otimes F$$

is independent of the basis  $B$ .

# Affine wreath product category

Start with the wreath product category  $\mathcal{W}(F)$ , but add a morphism:

$$\begin{array}{c} \uparrow \\ \circ \\ | \end{array} : \uparrow \rightarrow \uparrow$$

We impose the additional relations:

$$\begin{array}{c} \uparrow \\ \circ \\ \diagup \quad \diagdown \\ \diagdown \quad \diagup \\ \uparrow \end{array} - \begin{array}{c} \uparrow \\ \circ \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \uparrow \end{array} = \sum_{b \in B} \begin{array}{c} \uparrow \\ \bullet \\ | \end{array} \begin{array}{c} \uparrow \\ \bullet \\ | \end{array} b^{\vee} \quad , \quad \begin{array}{c} \uparrow \\ \bullet \\ | \\ \circ \\ | \end{array} = \begin{array}{c} \uparrow \\ \circ \\ | \\ \bullet \\ | \end{array} f \quad , \quad f \in F$$

Call the resulting category  $\mathcal{AW}(F)$  the **affine wreath product category**.

Now

$$\text{End}_{\mathcal{AW}(F)}(\uparrow^{\otimes n})$$

is an **affine wreath product algebra**.

**Note:**  $\mathcal{AW}(\mathbb{k})$  is the degenerate affine Hecke category.

# Adjunction

Suppose a strict monoidal category  $\mathcal{C}$  has two objects  $\uparrow$  and  $\downarrow$ , with

$$\text{id}_{\uparrow} = \uparrow \quad , \quad \text{id}_{\downarrow} = \downarrow .$$

A morphism  $\mathbf{1} \rightarrow \downarrow \otimes \uparrow$  would have string diagram

$$\begin{array}{c} \swarrow \quad \searrow \\ \vdots \end{array} \quad , \quad \text{where} \quad \begin{array}{c} \vdots \\ \vdots \end{array} = \text{id}_{\mathbf{1}} .$$

We typically omit the dotted line and draw:

$$\begin{array}{c} \swarrow \quad \searrow \end{array} : \mathbf{1} \rightarrow \downarrow \otimes \uparrow .$$

Similarly, we can have

$$\begin{array}{c} \searrow \quad \swarrow \end{array} : \uparrow \otimes \downarrow \rightarrow \mathbf{1} .$$

# Adjunction

We say that  $\downarrow$  is **right adjoint** to  $\uparrow$  (and  $\uparrow$  is **left adjoint** to  $\downarrow$ ) if there exist morphisms

$$\cup : \mathbf{1} \rightarrow \downarrow \otimes \uparrow \quad \text{and} \quad \cap : \uparrow \otimes \downarrow \rightarrow \mathbf{1}.$$

such that

$$\downarrow \cup = \downarrow \quad \text{and} \quad \cap \uparrow = \uparrow.$$

(This is analogous to the unit-counit formulation of adjunction of functors.)

We say  $\uparrow$  and  $\downarrow$  are **biadjoint** if they are both left and right adjoint to each other. So we also have

$$\cup : \mathbf{1} \rightarrow \uparrow \otimes \downarrow \quad \text{and} \quad \cap : \downarrow \otimes \uparrow \rightarrow \mathbf{1}$$

such that

$$\cup \uparrow = \uparrow \quad \text{and} \quad \cap \downarrow = \downarrow.$$

# The Frobenius Heisenberg category

Recall the affine wreath product category  $\mathcal{AW}(F)$ . It is the strict  $\mathbb{k}$ -linear monoidal category with:

**Objects:** Generated by object  $\uparrow$ .

**Morphisms:** Generated by

$$\begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} : \uparrow \otimes \uparrow \rightarrow \uparrow \otimes \uparrow, \\ \begin{array}{c} \uparrow \\ \circ \\ \uparrow \end{array} : \uparrow \rightarrow \uparrow, \quad \begin{array}{c} \uparrow \\ \bullet \\ \uparrow \end{array} f : \uparrow \rightarrow \uparrow, \quad f \in F,$$

with relations

$$\begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} = \uparrow \uparrow, \quad \begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} = \begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array}, \quad \begin{array}{c} \uparrow \\ \bullet \\ \circ \\ \uparrow \end{array} f = \begin{array}{c} \uparrow \\ \circ \\ \bullet \\ \uparrow \end{array} f, \quad f \in F, \\ \begin{array}{c} \uparrow \\ \circ \\ \nearrow \\ \searrow \end{array} - \begin{array}{c} \nearrow \\ \searrow \\ \circ \\ \uparrow \end{array} = \sum_{b \in B} \begin{array}{c} \uparrow \\ \bullet \\ \uparrow \end{array} b \begin{array}{c} \uparrow \\ \bullet \\ \uparrow \end{array} b^\vee, \quad \begin{array}{c} \uparrow \\ \bullet \\ \nearrow \\ \searrow \end{array} f = \begin{array}{c} \nearrow \\ \searrow \\ \bullet \\ \uparrow \end{array} f, \quad f \in F.$$

For  $n \in \mathbb{N}$ , define

$$\begin{array}{c} \uparrow \\ \circ \\ \uparrow \end{array} = \left. \begin{array}{c} \circ \\ \circ \\ \circ \\ \circ \\ \circ \end{array} \right\} n \text{ dots.}$$

# The Frobenius Heisenberg category

Fix a **central charge**  $\xi \in \mathbb{Z}$ ,  $\xi \leq 0$ .

(Actually, we can take any  $\xi \in \mathbb{Z}$ , but we choose  $\xi \leq 0$  for simplicity of exposition.)

To  $\mathcal{AW}(F)$  we add another object  $\downarrow$  that is **right adjoint** to  $\uparrow$ :

$$\begin{array}{c} \downarrow \\ \downarrow \end{array} = \downarrow \quad \text{and} \quad \begin{array}{c} \uparrow \\ \uparrow \end{array} = \uparrow.$$

We can then define **right crossings**:

$$\begin{array}{c} \nearrow \\ \searrow \\ \swarrow \\ \searrow \end{array} := \begin{array}{c} \uparrow \\ \downarrow \end{array} \begin{array}{c} \downarrow \\ \uparrow \end{array} : \uparrow\downarrow \rightarrow \downarrow\uparrow.$$

(We start denoting tensor product by juxtaposition:  $\uparrow\downarrow := \uparrow \otimes \downarrow$ .)

# The Frobenius Heisenberg category

We then impose the crucial **inversion relation**:

The following matrix of morphisms is an isomorphism in the additive envelope:

$$\left[ \begin{array}{c} \text{X} \\ \text{U} \end{array}, 0 \leq r \leq -\xi - 1, b \in B \right] : (\uparrow \otimes \downarrow) \oplus \mathbf{1}^{\oplus(-\xi \dim F)} \rightarrow \downarrow \otimes \uparrow .$$

More precisely, we add in some other morphisms that are the matrix components of an inverse to the above morphism.

We call the resulting category  $\mathcal{H}eis_{F,\xi}$  the **Frobenius Heisenberg category**.

# The Frobenius Heisenberg category

## Theorem (S. 2018)

There are unique morphisms

$$\cup : \mathbf{1} \rightarrow \uparrow\downarrow, \quad \cap : \downarrow\uparrow \rightarrow \mathbf{1} \quad (1)$$

such that the following relations hold:

$$\begin{aligned} \text{crossing} &= \uparrow \downarrow, & \text{crossing} &= \downarrow \uparrow + \sum_{k,s \geq 0} \sum_{a,b \in B} \text{diagram} \\ \text{loop} &= \delta_{\xi,0} \uparrow, & \text{loop} &= \delta_{r,-\xi-1} \text{tr}(f) \text{ if } 0 \leq r < -\xi. \end{aligned}$$

In addition  $\mathcal{H}eis_{F,\xi}$  can be presented equivalently by replacing the inversion relation with the existence of morphisms (1) and above relations.



# The Frobenius Heisenberg category

The previous theorem involves **left crossings**

$$\begin{array}{c} \nearrow \\ \searrow \\ \nwarrow \\ \nearrow \end{array} := \begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array}$$

and **negatively dotted** bubbles: for  $r \leq -\xi$ ,

$$r+\xi-1 \circlearrowleft \bullet f := (-1)^{r+1} \sum_{b_1, \dots, b_{r-1} \in B} \det \left( b_{j-1}^\vee b_j \circlearrowleft i-j-\xi \right)_{i,j=1}^r.$$

## Theorem (S. 2018)

- 1 The objects  $\downarrow$  and  $\uparrow$  are **biadjoint**.
- 2 The category  $\mathcal{H}eis_{F,\xi}$  is **strictly pivotal** (isotopy invariance for morphisms).
- 3 One can compute an infinite grassmannian relation, curl relations, bubble slide relations, and an alternating braid relation (omitted here).
- 4 Under a mild assumption on  $F$ , the category  $\mathcal{H}eis_{F,\xi}$  **categorifies** the Heisenberg algebra at central charge  $\xi \dim F$ .

## Heisenberg categorification and actions ( $\xi \leq -1$ )

Suppose  $\xi \leq -1$ .

The category  $\mathcal{H}eis_{F,\xi}$  acts naturally on modules for **cyclotomic wreath product algebras**. We have a chain of algebras

$$\mathbb{k} = A_0 \subseteq A_1 \subseteq A_2 \subseteq \cdots .$$

Then

- $\uparrow$  acts by induction from  $A_n\text{-mod}$  to  $A_{n+1}\text{-mod}$ ,
- $\downarrow$  acts by restriction from  $A_n\text{-mod}$  to  $A_{n-1}\text{-mod}$ .

The morphisms (diagrams) act by certain natural transformations.

Fact that  $\uparrow$  and  $\downarrow$  are biadjoint corresponds to fact that induction and restriction are biadjoint.

In other words  $A_n$  is a **Frobenius extension** of  $A_{n-1}$ .

# Heisenberg categorification and actions ( $\xi = 0$ )

## $F = \mathbb{k}$ case

$\mathcal{H}eis_{\mathbb{k},0}$  is the **affine oriented Brauer category** of Brundan–Comes–Nash–Reynolds.

$\mathcal{H}eis_{\mathbb{k},0}$  acts naturally on  $\mathfrak{gl}_n(\mathbb{k})$ -mod: If  $V$  is the natural rep, then

- $\uparrow \mapsto V \otimes -$
- $\downarrow \mapsto V^* \otimes -$

## General case: open problem

What does  $\mathcal{H}eis_{F,0}$  act naturally on for a general Frobenius algebra  $F$ ?

# Historical remarks

## Original Heisenberg category (Khovanov)

- Morphisms were planar diagrams **up to isotopy**, so strictly pivotal property was part of the definition.
- Central charge  $\xi = -1$  and  $F = \mathbb{k}$ .

## Frobenius modification (central charge $-1$ )

- For  $F$  the zigzag algebra, defined by Cautis–Licata and studied in relation to geometry of the **Hilbert scheme**.
- General definition given in joint work with Rosso.
- Still have central charge  $\xi = -1$ .

## Higher central charge (Mackaay–S.)

- Generalized to higher central charge (with  $F = \mathbb{k}$ ).
- Again, pivotal property part of the definition.

# Historical remarks

## Inversion relation approach, $F = \mathbb{k}$ (Brundan)


- New approach to the definition of higher charge category (Mackaay-S.) using the inversion relation.
- Now, pivotal property is a **consequence** of the definition.
- **Advantage**: proof that category acts on modules over degenerate (cyclotomic) affine Hecke algebras is much easier. Uses a well-known Mackey-type theorem.

## Current work

- Follows inversion relation approach of Brundan.
- Defines a Frobenius algebra version of higher charge category (Mackaay-S.).
- Defines a higher charge version of previous Frobenius Heisenberg category (Rosso-S.).

# Historical remarks

Summarizing the relationship between the Heisenberg categories appearing in the literature, we have:



## Some remarks

One can actually work in a more general setting than the one described here:

- 1  $F$  can be a **graded Frobenius superalgebra**. Then  $\mathcal{H}eis_{F,\xi}$  is a strict  $\mathbb{k}$ -linear **graded monoidal supercategory**.
- 2 The trace need not be symmetric. In general, there exists a **Nakayama automorphism**  $\psi: F \rightarrow F$  such that

$$\mathrm{tr}(fg) = (-1)^{\bar{f}\bar{g}} \mathrm{tr}(g\psi(f)) \quad \text{for all } f, g \in F.$$

Then, for instance,

$$f \begin{array}{c} \uparrow \\ \bullet \\ \circ \end{array} = \begin{array}{c} \uparrow \\ \circ \\ \bullet \end{array} \psi(f), \quad f \in F,$$

- 3 Above remarks mean we can take  $F$  to be the **Clifford superalgebra**. Then  $\mathcal{H}eis_{F,\xi}$  acts on modules for **affine Sergeev algebras** (a.k.a. **degenerate affine Hecke–Clifford algebras**).


# Work in progress (with J. Brundan)

One can  $q$ -deform the Frobenius Heisenberg category. When  $F = \mathbb{k}$ , this corresponds to



deg. affine Hecke algebra  $\rightsquigarrow$  affine Hecke algebra.

Generating objects:  $\uparrow$  and  $\downarrow$

Generating morphisms:



Relations: Fix  $z \in \mathbb{k}$ .



+ inversion relation.



## Work in progress (with J. Brundan)

Case:  $\xi = 0$

When  $\xi = 0$ , one obtains the **affine oriented skein category**, an affinization of the **HOMFLY-PT skein category**.

This category acts on modules for  $U_q(\mathfrak{gl}_n)$ .

Certain closed diagrams correspond to the **Casimir elements** in  $U_q(\mathfrak{gl}_n)$ .

Relation to previous constructions

When  $\xi = -1$ , the category contains the previously defined  **$q$ -deformed Heisenberg category** (Licata–S. 2013).

Main difference between two constructions is that, in the previous  $q$ -deformed Heisenberg category, the dot was not invertible.

Case  $\xi \neq 0$ : Action on modules for **cyclotomic Hecke algebras**.

## Work in progress (with J. Brundan)

Generally, one can again incorporate a graded Frobenius superalgebra to get a more general **quantum Frobenius Heisenberg category**.

When  $\xi \neq 0$ , category should act on cyclotomic quotients of quantum affine wreath product algebras. The theory of these algebras is yet to be developed.

When  $\xi = 0$ , the natural action is an open question for general  $F$ . Should be some  $F$ -deformation of  $U_q(\mathfrak{gl}_n)$ .