



**Part A: Answer Only Questions**

For Questions 1–13, only your final answer will be considered for marks.

1. **(1 point)** Which of the following statements is true for every matrix  $M$  with  $m$  rows and  $n$  columns, where  $m < n$ ?

- (a) The system  $M\mathbf{x} = \mathbf{0}$  has no solution.
- (b) The system  $M^T\mathbf{x} = \mathbf{0}$  has no solution.
- (c) The system  $M\mathbf{x} = \mathbf{0}$  has infinitely many solutions.
- (d) For every  $\mathbf{b}$  in  $\mathbb{R}^m$ , the system  $M\mathbf{x} = \mathbf{b}$  has infinitely many solutions.
- (e) For some  $\mathbf{b}$  in  $\mathbb{R}^m$ , the system  $M\mathbf{x} = \mathbf{b}$  has a unique solution.
- (f) For every  $\mathbf{b}$  in  $\mathbb{R}^n$ , the system  $M^T\mathbf{x} = \mathbf{b}$  has at least one solution.

**Answer:** \_\_\_\_\_

2. **(1 point)** Which of the following identities is **not** always true for  $n \times n$  matrices  $M$  and  $N$ ?

- (a)  $M + N = N + M$ ,
- (b)  $c(M + N) = cM + cN$ ,
- (c)  $(c + d)M = cM + dM$ ,
- (d)  $MN = NM$ ,
- (e)  $c(dM) = (cd)M$ ,
- (f)  $c(MN) = (cM)N$ .

**Answer:** \_\_\_\_\_

3. **(2 points)** Which of the following sets are subspaces of  $\mathbb{R}^3$ ? More than one answer may be correct.

- (a)  $\{(0, 0, 0)\}$
- (b)  $\{(a, b, 2) \mid a, b \in \mathbb{R}\}$
- (c)  $\{(a^2, 0, 0) \mid a \in \mathbb{R}\}$
- (d)  $\{(a^3, 0, b) \mid a, b \in \mathbb{R}\}$
- (e)  $\{(a, b, c) \mid a, b, c \in \mathbb{R}, a + 2b + 3c = 0\}$
- (f)  $\{(a, b, c) \mid a, b, c \in \mathbb{R}, a + 2b + 3c = 1\}$

**Answer:** \_\_\_\_\_

4. (1 point) Of the following matrices, circle those which are in reduced row echelon form.

$$\begin{array}{ccc} \begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, & \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, & \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \\ \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{bmatrix}, & \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}, & \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \\ \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, & \begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, & \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{bmatrix}. \end{array}$$

5. (3 points) If

$$M = \begin{bmatrix} 1 & 3 & 2 \\ 2 & -1 & 0 \\ -3 & 2 & -2 \end{bmatrix}, \quad \text{and} \quad N = \begin{bmatrix} 1 & -2 \\ 2 & 1 \\ 1 & 3 \end{bmatrix},$$

compute  $MN$ .

**Answer:**  $MN =$

6. (1 point) Compute  $(2 - 3i)(-1 + i)$ . Write your answer in the form  $a + bi$  where  $a$  and  $b$  are real numbers.

**Answer:** \_\_\_\_\_

Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

7. (2 points) Find real numbers  $a$  and  $b$  such that

$$a + bi = \frac{1 + 2i}{1 - i}.$$

Answer:  $a =$  \_\_\_\_\_  $b =$  \_\_\_\_\_

8. (1 point) Suppose  $\begin{vmatrix} a & c \\ b & d \end{vmatrix} = -2$ . What is  $\begin{vmatrix} 2b & 2d \\ -2a & -2c \end{vmatrix}$ ?

Answer: \_\_\_\_\_

9. (1 point) Let

$$M = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 2 & -1 & 0 & 0 \\ -1 & 4 & 2 & 0 \\ 3 & -2 & -1 & -2 \end{bmatrix} \quad \text{and} \quad N = \begin{bmatrix} 2 & 3 & -1 & 1 \\ 0 & 1 & 4 & 1 \\ 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$$

What is  $\det(MN)$ ?

Answer: \_\_\_\_\_

10. (1 point) Suppose

$$M = \begin{bmatrix} 1 & -6 & 8 & 10 & -5 & 12 & 0 \\ 0 & 0 & 4 & 5 & 7 & -3 & -2 \\ 0 & 0 & 0 & -2 & 0 & 8 & -5 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

What is the rank of  $M$ ?

Answer: \_\_\_\_\_

11. (1 point) For the matrix  $M$  of the previous question, what is the dimension of the null space of  $M$ ?

Answer: \_\_\_\_\_

12. (2 points) Suppose  $A$ ,  $B$  and  $C$  are  $6 \times 6$  matrices with

$$\det A = 2, \quad \det B = 7, \quad \det C = -1.$$

What is  $\det(-AB^3CC^T A^T B^{-3})$ ?

Answer: \_\_\_\_\_

13. (1 point) What are the eigenvalues of the matrix


$$\begin{bmatrix} 5 & 8 & 20 & -2 \\ 0 & -1 & 8 & 0 \\ 0 & 0 & 0 & 11 \\ 0 & 0 & 0 & 2 \end{bmatrix} ?$$

Answer: \_\_\_\_\_

**Part B: Long Answer Questions**

For Questions 14–22, you must show your work and justify your answers to receive full marks. Partial marks may be awarded for making sufficient progress towards a solution.

14. (3 points) Write the linear system describing the following traffic problem, labeling each equation by the intersection ( $A$ ,  $B$ ,  $C$ , or  $D$ ) to which it corresponds. **Do not solve the system.**



15. (6 points) Suppose the augmented matrix of a linear system is

$$\left[ \begin{array}{ccc|c} 1 & k-1 & 1 & 0 \\ 0 & 1 & k+1 & 1 \\ 0 & 0 & k^2-1 & k-1 \end{array} \right].$$

Determine the value(s) of  $k$  for which this system has

- (a) no solution,
- (b) infinitely many solutions,
- (c) a unique solution.

**Work / Justification:**

**Final Answer:** a) \_\_\_\_\_ b) \_\_\_\_\_ c) \_\_\_\_\_

Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

16. **(5 points)** Are the vectors  $(1, 1, 2)$ ,  $(1, 2, -3)$ , and  $(-1, 0, -7)$  linearly independent? If not, find a non-trivial linear combination of these three vectors that is equal to  $\mathbf{0}$ .



Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

17. (4 points) Find the inverse of the matrix

$$M = \begin{bmatrix} 2 & 7 & 1 \\ 1 & 4 & -1 \\ 1 & 3 & 0 \end{bmatrix}.$$

Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

18. (5 points) Suppose an economy with two sectors has unit consumption matrix

$$C = \begin{bmatrix} 0.1 & 0.4 \\ 0.3 & 0.2 \end{bmatrix}$$

and final demand vector

$$\mathbf{d} = \begin{bmatrix} 15 \\ 12 \end{bmatrix}.$$

Determine the production levels necessary to satisfy the given final demand. (Hint: use the inverse matrix method instead of row reduction.)

Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

19. (6 points) Find the eigenvalues of the matrix

$$M = \begin{bmatrix} 7 & -1 \\ 4 & 2 \end{bmatrix}.$$

For each eigenvalue, find the corresponding eigenspace.

20. (3 points) The eigenvalues of the matrix

$$M = \begin{bmatrix} -1 & 3 & -1 \\ -3 & 5 & -1 \\ -3 & 3 & 1 \end{bmatrix}$$

are 1 and 2. The eigenspace corresponding to the eigenvalue 1 is

$$\left\{ \begin{bmatrix} t \\ t \\ t \end{bmatrix}, t \in \mathbb{R} \right\},$$

and the eigenspace corresponding to the eigenvalue 2 is

$$\left\{ \begin{bmatrix} r \\ s \\ -3r + 3s \end{bmatrix}, r, s \in \mathbb{R} \right\}.$$

Find a matrix  $P$  such that the matrix  $P^{-1}MP$  is diagonal.

Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

21. (3 points) Let  $P$  and  $M$  be  $2 \times 2$  matrices with

$$P = \begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix}.$$

If

$$P^{-1}MP = \begin{bmatrix} 2 & 0 \\ 0 & -1 \end{bmatrix},$$

find  $M^7$ . Note: You do not need to compute powers of scalars (that is, you can leave expressions like  $2^n$  in your answer).

22. In a certain country, the mobile phone industry is dominated by two companies: Ten-Fours and Siren. Ten-Fours has 180 000 customers and Siren has 120 000 customers. Assume that, every year, 10% of the customer base of Ten-Fours switches to Siren and 5% of the customer base of Siren switches to Ten-Fours. For the purposes of this question, suppose no customer leaves a company without switching to the other one and no company attracts customers that are not leaving the other (that is, the only changes in customer base come from switching between the two companies).

(a). **(1 point)** Write down the transition (migration) matrix  $M$  and initial state vector  $\mathbf{x}_0$  for this problem.

(b). **(2 points)** Find the number of customers of Ten-Fours after one year.

(c). **(4 points)** Find the number of customers of Ten-Fours after many years. That is, find the long term stable number of customers of Ten-Fours.

Student # \_\_\_\_\_

MAT 1302A Final Exam, April 15, 2008

This page is for scrap work and is intentionally left blank.