


A graphical calculus for the Jack inner product on symmetric functions



Alistair Savage
University of Ottawa

Joint with: Anthony Licata (Australian National University) and Daniele Rosso (Indiana University Northwest)

Slides available online: alistairsavage.ca/talks

Paper: [arXiv:1610.01862](https://arxiv.org/abs/1610.01862) (to appear in *J. Comb. Theory A*)

Outline

Goal: Develop a graphical realization of the Jack inner product on symmetric functions.

Overview:

- 1 Review of symmetric functions & bilinear forms
- 2 Heisenberg algebras
- 3 Heisenberg categories
- 4 Planar diagrammatics: closed and annular diagrams
- 5 Diagrammatic bilinear form
- 6 Further directions

Power-sum and monomial symmetric functions

Let Sym be the \mathbb{C} -algebra of symmetric functions.

For $n \geq 1$, let

$$p_n = x_1^n + x_2^n + x_3^n + \cdots$$

be the n -th power sum.

For a partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell)$, define the power-sum symmetric function

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_\ell}.$$

We also define the monomial symmetric function

$$m_\lambda = \sum_{\alpha} x_1^{\alpha_1} x_2^{\alpha_2} \cdots,$$

where the sum is over all permutations $\alpha = (\alpha_1, \alpha_2, \dots)$ of λ .

Jack inner products

Fact: The p_λ form a basis of Sym .

Result: We can define an inner product on Sym by specifying its values on the p_λ .

Jack inner product

Fix $\alpha \in \mathbb{R}$, $\alpha \neq 0$. Define the **Jack inner product** by

$$\langle p_\lambda, p_\mu \rangle_\alpha = \delta_{\lambda, \mu} \alpha^{\ell(\lambda)} z_\lambda,$$

where $\ell(\lambda)$ is the length of λ and

$$z_\lambda = \prod_{k \geq 1} k^{m_k(\lambda)} m_k(\lambda)!,$$

where $m_k(\lambda)$ is the number of parts of λ equal to k .

Aside: Form is uniquely determined by its values on the p_n and the fact that it is a Hopf pairing.

Jack symmetric functions

Fact: The m_λ form a basis of Sym .

Dominance order: For partitions λ, μ we say

$$\mu \leq \lambda \iff \mu_1 + \cdots + \mu_i \leq \lambda_1 + \cdots + \lambda_i \text{ for all } i \geq 1.$$

Jack symmetric functions

The **Jack symmetric functions** J_λ are uniquely defined by the conditions

- 1 $J_\lambda \in m_\lambda + \text{Span}\{m_\mu \mid \mu < \lambda\}$,
- 2 $\langle J_\lambda, J_\mu \rangle_\alpha = 0$ if $\lambda \neq \mu$.

Aside: The Jack functions are obtained from the **Macdonald symmetric functions** via the specialization

$$q = t^\alpha, \quad t \rightarrow 1.$$

When $\alpha = 1$, the Jack functions are scalar multiples of the Schur functions.

The Heisenberg algebra

For the remainder of the talk, we fix $\alpha \in \mathbb{Z}$, $\alpha \neq 0$.

The Heisenberg algebra \mathfrak{h}_α

Let \mathfrak{h}_α be the \mathbb{C} -algebra generated by p_n^\pm , $n \in \mathbb{N}_+$, with relations

$$p_n^+ p_m^+ = p_m^+ p_n^+, \quad p_n^- p_m^- = p_m^- p_n^-, \quad p_n^- p_m^+ = p_m^+ p_n^- + \delta_{n,m} n \alpha.$$

As a vector space, we have

$$\mathfrak{h}_\alpha = \mathfrak{h}_\alpha^+ \otimes \mathfrak{h}_\alpha^-,$$

where

$$\mathfrak{h}_\alpha^\pm = \langle p_n^\pm \mid n \geq 1 \rangle \cong \text{Sym}$$

are subalgebras.

Action of the Heisenberg algebra on Sym

Action of \mathfrak{h}_α on Sym

\mathfrak{h}_α acts naturally on Sym:

- p_n^+ acts by multiplication by p_n ,
- p_n^- is adjoint to p_n^+ :

$$\langle p_n^- \cdot f, g \rangle_\alpha = \langle f, p_n g \rangle_\alpha \quad \forall f, g \in \text{Sym}.$$

This is called the **Fock space** representation of \mathfrak{h}_α .

Frobenius algebras

Frobenius algebras

A **graded Frobenius algebra** is a f.d. \mathbb{N} -graded algebra

$$B = \bigoplus_{n \in \mathbb{N}} B_n,$$

with a **trace map** $\text{tr}: B \rightarrow \mathbb{C}$ satisfying:

- tr is linear,
- the kernel of tr contains no nonzero left ideal.

Example

Let $B = \mathbb{C}[y]/(y^k)$, graded by degree.

Then B is a Frobenius algebra with trace map

$$\text{tr} \left(a_0 + a_1 y + \cdots + a_{k-1} y^{k-1} \right) = a_{k-1}.$$

The Heisenberg algebra \mathfrak{h}_B

For the rest of the talk, fix a graded Frobenius algebra

$$B = \bigoplus_{n=0}^{\delta} B_n, \quad B_{\delta} \neq 0.$$

Its **graded dimension** is

$$\text{grdim } B = \sum_{n \in \mathbb{N}} q^n \dim B_n \in \mathbb{N}[q].$$

Simplifying assumption (for this talk): $B_0 = \mathbb{C}$, $\delta > 0$.

The Heisenberg algebra \mathfrak{h}_B

Let \mathfrak{h}_B be the $\mathbb{C}(q)$ -algebra generated by p_n^{\pm} , $n \in \mathbb{N}_+$, with relations

$$p_n^+ p_m^+ = p_m^+ p_n^+, \quad p_n^- p_m^- = p_m^- p_n^-, \quad p_n^- p_m^+ = p_m^+ p_n^- + \delta_{n,m} n \text{ grdim } B.$$

Note: $\mathfrak{h}_B|_{q=1} = \mathfrak{h}_{\alpha}$, with $\alpha = \dim B$.

The Heisenberg category

We define a \mathbb{C} -linear strict monoidal category \mathcal{H}_B as follows:

The **objects** of \mathcal{H}_B are generated by two objects Q_+ and Q_- , together with their degree shifts.

So objects are sequences of Q_+ and Q_- , which a \mathbb{Z} -degree shift:

$$\emptyset, \quad \{2\}Q_+, \quad \{-4\}Q_-Q_-Q_+Q_+Q_+Q_-, \quad \text{etc.}$$

The **morphisms** are \mathbb{C} -linear combinations of planar diagrams:

- oriented compact one-manifolds immersed into the plane strip $\mathbb{R} \times [0, 1]$,
- agree with the domain/codomain at the top/bottom of the diagram,
- up to isotopy, and
- modulo certain local relations.

Morphisms: planar diagrams

We have the identity morphisms:

$$\text{id}_{Q_+} = \uparrow \qquad \text{id}_{Q_-} = \downarrow$$

Strands are allowed to carry dots labeled by elements of B :



Collision of dots is given by multiplication in B :

$$\begin{array}{c} \uparrow \\ \bullet \\ b \\ \bullet \\ b' \\ \uparrow \end{array} = \begin{array}{c} \uparrow \\ \bullet \\ b'b \\ \uparrow \end{array} \qquad \begin{array}{c} \downarrow \\ \bullet \\ b \\ \bullet \\ b' \\ \downarrow \end{array} = \begin{array}{c} \downarrow \\ \bullet \\ bb' \\ \downarrow \end{array}$$

Strands are allowed to cross:



Morphisms: grading

The morphism spaces are graded:


$$\text{deg } \begin{array}{c} \nearrow \\ \searrow \\ \swarrow \\ \searrow \end{array} = 0, \quad \text{deg } \begin{array}{c} \uparrow \\ \bullet \\ | \end{array} b = \text{deg } b,$$

$$\text{deg } \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \uparrow = 0, \quad \text{deg } \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \downarrow = 0,$$

$$\text{deg } \begin{array}{c} \uparrow \\ \text{---} \\ \text{---} \end{array} = \delta, \quad \text{deg } \begin{array}{c} \text{---} \\ \text{---} \\ \downarrow \end{array} = -\delta.$$

(Recall δ is the top degree of B .)

The degree of a morphism determines the relative degree shifts of the domain and codomain.




is a morphism

$$\{\text{deg } b_1 + \text{deg } b_2 + \text{deg } b_3 + \delta\} Q_- Q_+ Q_+ Q_+ Q_- Q_+ \rightarrow Q_+ Q_- Q_+ Q_+$$

Morphisms: local relations

The **local relations** we impose are:



Here \mathcal{B} is a basis of B and $\{b^v \mid b \in \mathcal{B}\}$ is the (right) dual basis:

$$\text{tr}(bc^v) = \delta_{b,c}.$$

Remark: Grothendieck group categorification

Recall: The **Karoubi envelope** (or **idempotent completion**) is an “enlargement” of a category \mathcal{C} , with one object for each idempotent morphism in \mathcal{C} .

Let $\text{Kar } \mathcal{H}_B$ be the Karoubi envelope of the graphical category \mathcal{H}_B .

Theorem (Rosso–S. 2015)

There is a natural algebra isomorphism

$$\mathfrak{h}_B \rightarrow \text{split Grothendieck group of } \text{Kar } \mathcal{H}_B.$$

Hence $\text{Kar } \mathcal{H}_B$ **categorifies** \mathfrak{h}_B .

Natural action of \mathcal{H}_B

For $n \geq 1$, consider the wreath product algebra

$$B^{\otimes n} \rtimes S_n.$$

As a vector space, $B^{\otimes n} \rtimes S_n = B^{\otimes n} \otimes \mathbb{C}S_n$.

The factors $B^{\otimes n}$ and $\mathbb{C}S_n$ are subalgebras, and

$$wb = (w \cdot b)w, \quad b \in B^{\otimes n}, w \in S_n,$$

where $w \cdot b$ is action of w on b by permuting factors.

Action of \mathcal{H}_B on $\bigoplus_n B^{\otimes n} \rtimes S_n$ -mod

- Q_+ acts by induction $B^{\otimes n} \rtimes S_n$ -mod $\rightarrow B^{\otimes(n+1)} \rtimes S_{n+1}$,
- Q_- acts by restriction $B^{\otimes(n+1)} \rtimes S_{n+1} \rightarrow B^{\otimes n} \rtimes S_n$ -mod,
- planar diagrams are natural transformations between compositions of induction and restriction.

Center of a monoidal category

Suppose \mathcal{C} is an additive monoidal category.

Then \mathcal{C} has an **identity object** $\mathbf{1}$:

$$X \otimes \mathbf{1} \cong X \cong \mathbf{1} \otimes X \quad \text{for all objects } X \text{ in } \mathcal{C}.$$

Center of a monoidal category

The **center** of \mathcal{C} is

$$Z(\mathcal{C}) := \text{End}_{\mathcal{C}}(\mathbf{1}).$$

The center is naturally a ring, under sum and tensor product of morphisms.

Example

If R is a ring and \mathcal{C} is the category of (R, R) -bimodules, then


$$Z(\mathcal{C}) = \text{End}_{\mathcal{C}}(R) \cong Z(R).$$

Closed diagrams

The identity object of \mathcal{H}_B is \emptyset .

So the center $Z(\mathcal{H}_B)$ consists of linear combinations of **closed diagrams**.

Multiplication is given by juxtaposition.



This algebra is clearly commutative (since we consider diagrams up to isotopy).

Trace of a monoidal category

Suppose \mathcal{C} is a \mathbb{C} -linear monoidal category.

The **trace** (or **zeroth Hochschild homology**) is

$$\mathrm{Tr}(\mathcal{C}) = \left(\bigoplus_{x \in \mathrm{Ob} \mathcal{C}} \mathrm{End}_{\mathcal{C}}(x) \right) / \mathrm{Span}\{fg - gf \mid f: x \rightarrow y, g: y \rightarrow x\}.$$

Example

Suppose A is an algebra. Let \mathcal{C} be the monoidal category with

- one object \star ,
- $\mathrm{Hom}_{\mathcal{C}}(\star, \star) = A$,
- tensor product of morphisms given by the product in A .


Then

$$\mathrm{Tr}(\mathcal{C}) = A / \mathrm{Span}\{ab - ba \mid a, b \in A\}$$

is the **trace** (or **cocenter**) of A .

Annular diagrams

The trace $\text{Tr}(\mathcal{H}_B)$ of \mathcal{H}_B can be identified the space of **annular diagrams**.






This is an algebra, with product given by **nesting**.

If A_1 and A_2 are annular diagrams, then A_1A_2 is the annular diagram obtained by placing A_2 in the center region of A_1 .


Diagrammatic action

The algebra $\text{Tr}(\mathcal{H}_B)$ of annular diagrams **acts** on the space $Z(\mathcal{H}_B)$ of closed diagrams.




Closures of permutations

Identify permutations with diagrams:





Given a permutation w , we can close off to the right to get a closed diagram or an annular diagram:



Key elements in the trace

More generally, any element of $\mathbb{C}S_n$ can be drawn as a linear combination of braid-like diagrams, on upward or downward strands.

Closing off to the right yields an annular diagram.



So to any $f \in \mathbb{C}S_n$, we have the associated annular diagram $[f^\pm]$. Let

$$P_\lambda^\pm = [w_\lambda^\pm],$$

where w_λ is a permutation of cycle type λ .

Question: Why is this well defined?

Example of a diagrammatic proof

Lemma

If $w_1, w_2 \in S_n$ are conjugate, then $[w_1^\pm] = [w_2^\pm]$.

Proof.

Suppose $w_1 = ww_2w^{-1}$. Then

$$[w_1^+] = \begin{array}{c} \text{---} \\ \uparrow \\ \boxed{w_1} \\ \downarrow \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \uparrow \\ \boxed{w^{-1}} \\ \uparrow \\ \boxed{w_2} \\ \uparrow \\ \boxed{w} \\ \downarrow \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \uparrow \\ \boxed{w_2} \\ \uparrow \\ \boxed{w} \\ \uparrow \\ \boxed{w^{-1}} \\ \downarrow \\ \text{---} \end{array} = \begin{array}{c} \text{---} \\ \uparrow \\ \boxed{w_2} \\ \downarrow \\ \text{---} \end{array} = [w_2^+].$$

□

Identification of the trace

Theorem (Licata–Rosso–S. 2016)

We have an isomorphism of algebras

$$\mathrm{Tr}(\mathcal{H}_B) \cong \mathfrak{h}_B, \quad P_\lambda^\pm \mapsto p_\lambda^\pm.$$

Note: We also have

$$[e_\lambda^\pm] \mapsto s_\lambda^\pm,$$

where $e_\lambda \in \mathbb{C}S_n$ is the primitive idempotent corresponding to a partition λ and s_λ is the **Schur function**.

The diagrammatic bilinear form

Let $\mathrm{Tr}(\mathcal{H}_B)^\pm$ be the algebra of annular diagrams spanned by closing off endomorphisms of $(\mathbb{Q}_\pm)^n$ for $n \in \mathbb{N}$.

So

$$\begin{aligned}\mathrm{Tr}(\mathcal{H}_B)^+ &= \text{algebra of clockwise annular diagrams,} \\ \mathrm{Tr}(\mathcal{H}_B)^- &= \text{algebra of counterclockwise annular diagrams.}\end{aligned}$$

Under the isomorphism $\mathrm{Tr}(\mathcal{H}_B) \cong \mathfrak{h}_B$, we have

$$\mathrm{Tr}(\mathcal{H}_B)^\pm \mapsto \mathfrak{h}_B^\pm := \langle p_n^\pm \mid n \in \mathbb{N} \rangle \cong \mathrm{Sym}.$$

So a pairing

$$\mathrm{Tr}(\mathcal{H}_B)^- \times \mathrm{Tr}(\mathcal{H}_B)^+ \rightarrow \mathbb{C}$$

corresponds to a **bilinear form** on Sym .

The diagrammatic bilinear form

Fact 1: The space $Z(\mathcal{H}_B)$ of closed diagrams is nonnegatively graded.

Fact 2: The only degree zero closed diagram is the empty diagram.

So we have the **projection onto degree 0**:

$$\mathbf{F}_0: Z(\mathcal{H}_B) \rightarrow \mathbb{C}.$$

The diagrammatic pairing

We define a pairing

$$\begin{aligned} \langle -, - \rangle_B: \operatorname{Tr}(\mathcal{H}_B)^- \times \operatorname{Tr}(\mathcal{H}_B)^+ &\rightarrow \mathbb{C}, \\ \langle x, y \rangle_B &= \mathbf{F}_0((xy) \cdot 1_\emptyset), \end{aligned}$$

where 1_\emptyset is the empty diagram.

The diagrammatic pairing

Graphically, $\langle x, y \rangle$ is obtained by

- placing the annular diagram y inside the annular diagram x ,
- viewing the resulting annular diagram as a closed diagram,
- projecting onto degree zero.

$$\langle x, y \rangle_B = \mathbf{F}_0 \left(\begin{array}{c} \text{Diagram of } x \text{ and } y \text{ nested} \end{array} \right) .$$

Note: If $f, g \in \mathbb{C}S_n$, then the nested diagram $[f][g] \cdot 1_\emptyset$ is already in degree zero and \mathbf{F}_0 is unnecessary.

The diagrammatic pairing

Theorem (Licata–Rosso–S.)

Under the isomorphisms $\mathrm{Tr}(\mathcal{H}_B)^\pm \cong \mathrm{Sym}$, the diagrammatic form corresponds to the Jack bilinear form at parameter

$$\alpha = \dim B.$$

So we have a **categorification** of the Jack bilinear form, with the Jack parameter categorified by the graded Frobenius algebra B .

Example

Suppose $B = \mathbb{C}[y]/(y^k)$. Trace map is the coefficient of y^{k-1} .

The first power sum p_1 corresponds to the clockwise circle P_1^+ and counterclockwise circle P_1^- .

$$\begin{aligned} \langle P_1^-, P_1^+ \rangle_B &= \text{diagram of two concentric circles} = \text{diagram of two overlapping circles} + \sum_{j=0}^{k-1} \text{diagram of a circle with a hole and two blue dots} \\ &= \text{diagram of two separate circles} + \sum_{j=0}^{k-1} y^{k-j-1} \text{diagram of a circle with a blue dot} = k. \end{aligned}$$

This agrees with the inner product of p_1 with itself in the Jack inner product at Jack parameter $k = \dim B$.

Full generality

The full results are more general than presented in this talk.

The Frobenius algebra B

- In general, B can be a graded Frobenius **superalgebra**.
- We don't need $B_0 = \mathbb{C}$.
- We only assume that all simple B -modules are of **type M** (i.e. not isomorphic to their parity shifts) and that the trace map is supersymmetric and even.
- The Jack parameter α corresponds to $\dim B_{\text{even}} - \dim B_{\text{odd}}$.

Further directions I

More general Frobenius algebras

- The categories \mathcal{H}_B are defined for an **arbitrary** graded Frobenius superalgebra B .
- Allowing B to have simple modules of type Q (isomorphic to their own parity shifts) would result in the space of **Schur Q -functions**.
- Allowing the trace map to not be supersymmetric would introduce **twisted Heisenberg algebras**.

Connections to W -algebras

- $\mathrm{Tr}(\mathcal{H}_{\mathbb{C}})$ is isomorphic to a quotient of the **W -algebra** $W_{1+\infty}$ (Cautis–Licata–Lauda–Sussan)
- $\mathrm{Tr}(\mathcal{H}_B)$ should be related to W -algebras associated to the lattice $K_0(B\text{-mod})$.

Further directions II

Wreath product algebras

- \mathcal{H}_B acts on modules for wreath product algebras $B^{\otimes n} \rtimes S_n$.
- Thus, the Heisenberg algebra $\mathrm{Tr}(\mathcal{H}_B) \cong \mathfrak{h}_B$ acts on the centers of these module categories.
- Can therefore use diagrammatics to study the centers of these categories.

Jack symmetric functions

- We have categorified the Jack inner product.
- **Question:** Can we categorify the Jack symmetric functions themselves?
- Find natural annular diagrams that correspond to these functions.

Further directions III

Geometry: Hilbert schemes

- Equivariant K -theory of the **Hilbert scheme** of points on \mathbb{C}^2 is related to the Macdonald ring of symmetric functions (Haiman).
- Equivariant homology related to Jack symmetric functions (Nakajima, Li–Qin–Wang).
- “ K -theory versus homology” is analogous to “Grothendieck group versus trace”.
- So current work should be related to these geometric constructions.